

# Tropical Boubou

*Laniarius aethiopicus*

1,199 plots from 3,488 records

Britton (1980) gives 4 races for Tanzania. *L.a. sublacteus* throughout the coastal lowlands inland to Mpwapwa and the North Pares. The highland race *L.a. ambiguus* south to Mts Meru, Kilimanjaro and Monduli. *L.a. major* in the W and then E to Crater Highlands, Njombe and Mahenge. *L.a. mossambicus* in the SE, S of *major* and *sublacteus*.

Localities mentioned in Britton.  
*sublacteus* - red circle  
*ambiguus* - red square  
*major* - blue circle

That gap through sukumaland, into the western Miombo and through the Bahi depression to the Masai Steppe is most probably real.

Virgin Yellow-vented Bulbul squares **12**

Single YvB month **21**

Two YvB months **24**

### Seasonality

Month	presence	bs	eggs
January	365	9	3
February	285	12	2
March	267	15	3
April	202	5	-
May	215	7	3
June	267	6	2
July	333	5	1
August	289	4	2
September	271	10	2
October	316	5	2
November	302	9	-
December	251	9	1
month 13	8		


### Status

- Presence
- Breeding season
- Month with eggs

- 850 - 900
- 900 - 950
- 950 - 1,000
- 1,000 - 1,050
- 1,050 - 1,100
- 1,100 - 1,200
- 1,200 - 1,300
- 1,300 - 1,400
- 1,400 - 1,500
- 1,500 - 1,600
- 1,600 - 1,700
- 1,700 - 1,800
- 1,800 - 1,900
- 1,900 - 2,000
- 2,000 - 2,100
- 2,100 - 2,200
- 2,200 - 2,300
- 2,300 - 2,400
- 2,400 - 2,500

### 30m SRTM

- 0 - 100
- 100 - 200
- 200 - 300
- 300 - 400
- 400 - 450
- 450 - 500
- 500 - 550
- 550 - 600
- 600 - 650
- 650 - 700
- 700 - 750
- 750 - 800
- 800 - 850



Tanzania Bird Atlas  
 Preliminary Map  
 September 2009  
 From 862,509  
 Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records  
 within each square