

Black-headed Gonolek

Laniarius erythrogaster

115 plots from 291 records

Virgin Yellow-vented Bulbul squares **12**

Single YvB month **21**

Two YvB months **24**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	32	1	-
February	32	-	-
March	22	-	-
April	13	-	-
May	13	1	-
June	27	-	-
July	62	3	-
August	18	-	-
September	11	-	-
October	14	-	-
November	17	1	-
December	21	-	-
month 13	3	-	-

Status

- Presence
- Breeding season
- Month with eggs

30m SRTM

- 0 - 100
- 100 - 200
- 200 - 300
- 300 - 400
- 400 - 450
- 450 - 500
- 500 - 550
- 550 - 600
- 600 - 650
- 650 - 700
- 700 - 750
- 750 - 800
- 800 - 850
- 850 - 900
- 900 - 950
- 950 - 1,000
- 1,000 - 1,050
- 1,050 - 1,100
- 1,100 - 1,200
- 1,200 - 1,300
- 1,300 - 1,400
- 1,400 - 1,500
- 1,500 - 1,600
- 1,600 - 1,700
- 1,700 - 1,800
- 1,800 - 1,900
- 1,900 - 2,000
- 2,000 - 2,100
- 2,100 - 2,200
- 2,200 - 2,300
- 2,300 - 2,400
- 2,400 - 2,500

Virtually confined to the Lake Victoria basin but those 2 records from Mwadui (3303D) where natural habitat is protected within the Williamson Diamonds property and those 2 records in the Mwowosi suggest a far larger range in the recent past. One of those latter records is a specimen in the Field Museum (Nobby to check) but the other is quite recent from the TGT consession so this bird may well still exist in numbers along the Malagarasi.

No previous records from 3101D so these birds that Bob Matthews is reporting from Bumbire are new for that square.

No records from Rubondo NP which is rather curious.

Tanzania Bird Atlas
Preliminary Map
September 2009
From 862,509
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records
within each square

